

Hydrogen selection rules

By computing transition elements, verify the following selection rules.

$$|l_b - l_a| = 1, \quad |m_b - m_a| \leq 1$$

Transition element.

$$T_{a \rightarrow b} = |\langle \psi_b | \hat{x} | \psi_a \rangle|^2 + |\langle \psi_b | \hat{y} | \psi_a \rangle|^2 + |\langle \psi_b | \hat{z} | \psi_a \rangle|^2$$

Transition amplitudes.

$$\langle \psi_b | \hat{x} | \psi_a \rangle = \int_V \psi_b^* \hat{x} \psi_a dV, \quad \langle \psi_b | \hat{y} | \psi_a \rangle = \int_V \psi_b^* \hat{y} \psi_a dV, \quad \langle \psi_b | \hat{z} | \psi_a \rangle = \int_V \psi_b^* \hat{z} \psi_a dV$$

Operators for dipole approximation.

$$\hat{x} = r \sin \theta \cos \phi, \quad \hat{y} = r \sin \theta \sin \phi, \quad \hat{z} = r \cos \theta$$

Volume measure for spherical coordinates.

$$dV = r^2 \sin \theta dr d\theta d\phi$$

Hydrogen wave functions ψ_{nlm} are solutions to the time-independent Schrödinger equation.

$$\hat{H} \psi_{nlm}(r, \theta, \phi) = E_n \psi_{nlm}(r, \theta, \phi)$$

Hydrogen wave function.

$$\psi_{nlm}(r, \theta, \phi) = R_{nl}(r) Y_{lm}(\theta, \phi)$$

Radial wave function.

$$R_{nl}(r) = \frac{2}{n^2} \sqrt{\frac{(n-l-1)!}{(n+l)!}} \left(\frac{2r}{na_0} \right)^l L_{n-l-1}^{2l+1} \left(\frac{2r}{na_0} \right) \exp \left(-\frac{r}{na_0} \right) a_0^{-3/2}$$

Associated Laguerre polynomial.

$$L_n^m(x) = (n+m)! \sum_{k=0}^n \frac{(-x)^k}{(n-k)!(m+k)!k!}$$

Spherical wave function.

$$Y_{lm}(\theta, \phi) = (-1)^m \sqrt{\frac{(2l+1)}{4\pi} \frac{(l-m)!}{(l+m)!}} P_l^m(\cos \theta) \exp(im\phi)$$

Associated Legendre polynomial of $\cos \theta$.

$$P_l^m(\cos \theta) = \begin{cases} \left(\frac{\sin \theta}{2} \right)^m \sum_{k=0}^{l-m} (-1)^k \frac{(l+m+k)!}{(l-m-k)!(m+k)!k!} \left(\frac{1-\cos \theta}{2} \right)^k, & m \geq 0 \\ (-1)^m \frac{(l+m)!}{(l-m)!} P_l^{|m|}(\cos \theta), & m < 0 \end{cases}$$