

Exercise 6.6. Assume Charlie has prepared the two spins in the singlet state. This time, Bob measures  $\tau_y$  and Alice measures  $\sigma_x$ . What is the expectation value of  $\sigma_x\tau_y$ ?

What does this say about the correlation between the two measurements?

---

Recall that

$$|sing\rangle = \frac{|ud\rangle - |du\rangle}{\sqrt{2}}$$

The expectation value of  $\sigma_x\tau_y$  is

$$\langle\sigma_x\tau_y\rangle = \langle sing|\sigma_x\tau_y|sing\rangle$$

Applying the  $\tau_y$  operator first we have from Table 1 the following result.

$$\langle\sigma_x\tau_y\rangle = \langle sing|\sigma_x\left(\frac{-i|uu\rangle - i|dd\rangle}{\sqrt{2}}\right)$$

Then applying the  $\sigma_x$  operator we have

$$\langle\sigma_x\tau_y\rangle = \langle sing|\left(\frac{-i|du\rangle - i|ud\rangle}{\sqrt{2}}\right)$$

Hence

$$\begin{aligned}\langle\sigma_x\tau_y\rangle &= \frac{1}{2}(\langle ud| - \langle du|)(-i|du\rangle - i|ud\rangle) \\ &= \frac{1}{2}(-i\langle ud|ud\rangle + i\langle du|du\rangle) \\ &= 0\end{aligned}\tag{1}$$

From (1) and the result  $\langle\sigma_x\rangle = \langle\tau_y\rangle = 0$  given on page 174 we have

$$\langle\sigma_x\tau_y\rangle - \langle\sigma_x\rangle\langle\tau_y\rangle = 0$$

Hence the measurements  $\sigma_x$  and  $\tau_y$  are uncorrelated.